

(11) Publication number: 2000106893 A

Generated Document.

PATENT ABSTRACTS OF JAPAN

(21) Application number: 10283845

(22) Application date: 06.10.98

(51) Intl. Cl.: C12P 17/02 C07D305/14 C12N 1/38

(30) Priority:

(43) Date of application publication:

18.04.00

(84) Designated contracting

states:

(71) Applicant: KOBE TENNENBUTSU KAGAKU

(72) Inventor: ANDO MASAYOSHI

(74) Representative:

(54) **PRODUCTION OF** TAXANE-TYPE DITERPENE

(57) Abstract:

PROBLEM TO BE SOLVED: To provide a method for producing a taxane-type diterpene such as taxol or the like, which can cope with increase in quantity demanded, can be carried out at a low cost and can preserve the environment.

SOLUTION: A taxane-type diterpene is took out by extracting, isolating and purifying callus prepared through a tissue culture method of needle leave of Taxus cuspidata Sieb et Zucc. or Taxus cuspidata Sieb et Zucc. var. nana Rehder which have been collected in the season when monthly. mean temperature for a month before being collected is ≤180°C. The culture medium includes at least one kind selected from a group consisting of 1naphthylacetic acid, 2,4dichlorophenoxyacetic acid, 4chloroindole-3-acetic acid, methyl jasmonate and an oligosaccharide more preferably.

COPYRIGHT: (C)2000,JPO

The Delphion Integrated View

Buy Now: More choices...

Tools: Add to Work File: Create new Work File

Go

View <u>Image</u>

1 page

View: Jump to: Top

Email this to a friend

學Title: JP2000106893A2: PRODUCTION OF TAXANE-TYPE DITERPENE

\$ Country:

JP Japan

ଟKind:

A2 Document Laid open to Public inspection i

☑ Inventor:

ANDO MASAYOSHI:

PAssignee:

KOBE TENNENBUTSU KAGAKU KK

News, Profiles, Stocks and More about this company

Published / Filed:

April 18, 2000 / Oct. 6, 1998

Number:

JP1998000283845

땋IPC Code:

C12P 17/02; C07D 305/14; C12N 1/38; C12P 17/02;

Priority Number:

Oct. 6, 1998 JP1998000283845

8 Abstract:

PROBLEM TO BE SOLVED: To provide a method for producing a taxane-type diterpene such as taxol or the like, which can cope with increase in quantity demanded, can be carried out at a low cost and can preserve the environment.

SOLUTION: A taxane-type diterpene is took out by extracting, isolating and purifying callus prepared through a tissue culture method of needle leave of Taxus cuspidata Sieb et Zucc. or Taxus cuspidata Sieb et Zucc. var. nana Rehder which have been collected in the season when monthly mean temperature for a month before being collected is ≤180°C. The culture medium includes at least one kind selected from a group consisting of 1naphthylacetic acid, 2,4-dichlorophenoxyacetic acid, 4-chloroindole-3-acetic acid, methyl jasmonate and an oligosaccharide more preferably.

COPYRIGHT: (C)2000,JPO

당 Family:

None

TOTAL Other Abstract

None









this for the Gallery...

© 1997-2002 Delphion, Inc.

Research Subscriptions | Privacy Policy | Terms & Conditions | Site Map | Contact Us